

NEW PHASE STRUCTURE OF THE NAMBU – JONA - LASINIO MODEL AT NONZERO CHEMICAL POTENTIAL

K.G. Klimenko[†], A.S. Vshivtsev[‡],

Institute for High Energy Physics, 142284, Protvino, Moscow region, Russia

E-mail: [†] kklm@mx.ihep.su, [‡] alexandr@vvas.msk.ru

It is shown that in the Nambu – Jona - Lasinio model at nonzero chemical potential there are two different massive phases with spontaneously broken chiral symmetry. In one of them particle density is identically zero, in another phase it is not equal to zero. The transition between phases is a phase transition of the second order.

This is an english translation of the article, published in Russian journal Pis'ma Zh. Eksp. Teor. Fiz. (JETP Letters) [1].

It is well known that Nambu – Jona - Lasinio (NJL) model [2, 3] is perfect laboratory for the investigations of low energy region in QCD. The subject of special interest is the spontaneous breaking of chiral symmetry. In the framework of NJL model this phenomenon was studied at nonzero temperature and density [4, 5], in the presence of external fields [6], at nonzero curvature and nontrivial topology of space - time [7].

In the present article the phase structure of NJL model at nonzero chemical potential μ is considered. In contrast with earlier papers [4, 5] we have found new massive phase of the model. The NJL Lagrangian has the form:

$$L = \sum_{k=1}^N \bar{\psi}_k i \hat{\partial} \psi_k + \frac{G}{2N} [(\sum_{k=1}^N \bar{\psi}_k \psi_k)^2 + (\sum_{k=1}^N \bar{\psi}_k i \gamma_5 \psi_k)^2]. \quad (1)$$

It contains N four - components Dirac fields ψ . So one can use $1/N$ - expansion technique. Eq. (1) is invariant under continuous chiral transformations:

$$\psi_k \rightarrow e^{i\theta\gamma_5} \psi_k (k = 1, \dots, N). \quad (2)$$

First of all let us recall some well - known vacuum properties of the theory (1) at $\mu = 0$. The introduction of an auxiliary Lagrangian

$$\tilde{L} = \bar{\psi} i \hat{\partial} \psi - \bar{\psi} (\sigma_1 + i \sigma_2 \gamma_5) \psi - \frac{N}{2G} (\sigma_1^2 + \sigma_2^2) \quad (3)$$

greatly facilitates the problem under consideration. (In (3) and other formulae below we have omitted the fermionic index k for simplicity.) On the equations of motion for auxiliary bosonic fields $\sigma_{1,2}$ the theory (3) is equivalent to the (1) one.

From (3) it follows in the leading order of $1/N$ - expansion:

$$\exp(iS_{eff}(\sigma_{1,2})) = \int D\bar{\psi} D\psi \exp(i \int \tilde{L} d^4x),$$

where

$$\frac{1}{N}S_{eff}(\sigma_{1,2}) = - \int d^4x \frac{\sigma_1 + \sigma_2}{2G} - i \ln \det(i\hat{\partial} - \sigma_1 - i\gamma_5\sigma_2). \quad (4)$$

Supposing that in this formula $\sigma_{1,2}$ are not dependent from the space - time points we have by definition:

$$S_{eff}(\sigma_{1,2}) = -V_{eff}(\sigma_{1,2}) \int d^4x, \quad (5)$$

where ($\Sigma = \sqrt{\sigma_1^2 + \sigma_2^2}$):

$$\frac{1}{N}V_{eff}(\sigma_{1,2}) = \frac{\Sigma^2}{2G} + 2i \int \frac{d^4p}{(2\pi)^4} \ln(\Sigma^2 - p^2) \equiv \frac{1}{N}V_0(\Sigma). \quad (6)$$

Introducing in (6) Euclidean metrics ($p_0 \rightarrow ip_0$) and cutting off the range of integration ($p^2 \leq \Lambda^2$), we obtain:

$$\begin{aligned} \frac{1}{N}V_0(\Sigma) &= \frac{\Sigma^2}{2G} - \frac{1}{16\pi^2} \left\{ \Lambda^4 \ln \left(1 + \frac{\Sigma^2}{\Lambda^2} \right) + \Lambda^2 \Sigma^2 - \right. \\ &\quad \left. - \Sigma^4 \ln \left(1 + \frac{\Lambda^2}{\Sigma^2} \right) \right\}. \end{aligned} \quad (7)$$

The stationary equation for the effective potential (7) has the form:

$$\frac{\partial V_0(\Sigma)}{\partial \Sigma} = 0 = \frac{N\Sigma}{4\pi^2} \left\{ \frac{4\pi^2}{G} - \Lambda^2 + \Sigma^2 \ln \left(1 + \frac{\Lambda^2}{\Sigma^2} \right) \right\} \equiv \frac{N\Sigma}{4\pi^2} F(\Sigma). \quad (8)$$

Now one can easily see that at $G < G_c = 4\pi^2/\Lambda^2$ eq. (8) has no solutions apart from $\Sigma = 0$. Hence, in this case fermions are massless, and chiral invariance (2) is not broken.

If $G > G_c$, then Eq. (8) has one nontrivial solution $\Sigma_0(G, \Lambda) \neq 0$ such that $F(\Sigma_0) = 0$. In this case Σ_0 is a point of global minimum for the potential $V_0(\Sigma)$. This means the spontaneous breaking of the symmetry (2) takes place. Moreover, fermions acquire mass $M \equiv \Sigma_0(G, \Lambda)$.

In the following at $G > G_c$ we shall use the fermionic mass M as an independent parameter of the theory. Three quantities G , M and Λ are connected by the Eq. (8). In terms of M and Λ the effective potential $V_0(\Sigma)$ has an equivalent form:

$$\begin{aligned} \frac{16\pi^2}{N} V_0(\Sigma) = & \Sigma^2 \Lambda^2 - 2\Sigma^2 M^2 \ln(1 + \Lambda^2/M^2) - \\ & - \Lambda^4 \ln(1 + \Sigma^2/\Lambda^2) + \Sigma^4 \ln(1 + \Lambda^2/\Sigma^2) \end{aligned} \quad (9)$$

Let us now imagine that $\mu > 0$ and temperature $T \neq 0$. In this case one can obtain effective potential $V_{\mu T}(\Sigma)$ if the measure of integration in (6) is transformed as

$$\int \frac{dp_0}{2\pi} \rightarrow iT \sum_{n=-\infty}^{\infty}, \quad p_0 \rightarrow i\pi T(2n+1) + \mu.$$

Summing there over n [8], we have:

$$\begin{aligned} \frac{1}{N} V_{\mu T}(\Sigma) = & \frac{1}{N} V_0(\Sigma) - 2T \int \frac{d^3 p}{(2\pi)^3} \ln \{ [1 + \\ & + \exp(-\beta(\sqrt{\Sigma^2 + \vec{p}^2} + \mu))] [1 + \exp(-\beta(\sqrt{\Sigma^2 + \vec{p}^2} - \mu))] \}, \end{aligned} \quad (10)$$

where $\beta = 1/T$, and $V_0(\Sigma)$ is presented in (7-9). Since we are going to study phase structure of NJL model at $\mu > 0$ and $T = 0$ only, one should direct the temperature to the zero in the above equation. From this we have corresponding effective potential $V_\mu(\Sigma)$:

$$\begin{aligned} \frac{1}{N} V_\mu(\Sigma) = & \frac{1}{N} V_0(\Sigma) - \frac{\Theta(\mu - \Sigma)}{16\pi^2} \left\{ \frac{10}{3} \mu (\mu^2 - \Sigma^2)^{3/2} - \right. \\ & \left. - 2\mu^3 \sqrt{\mu^2 - \Sigma^2} + \Sigma^4 \ln [(\mu + \sqrt{\mu^2 - \Sigma^2})^2 / \Sigma^2] \right\}. \end{aligned} \quad (11)$$

The function (11) is symmetric under transformation $\Sigma \rightarrow -\Sigma$, so it is sufficient to learn the situation with $\Sigma \geq 0$ only.

In the present paper we shall consider the case $G > G_c$ only. From (9) and (11) one can find stationary equation for the effective potential (M, Λ are free parameters):

$$\frac{\partial V_\mu(\Sigma)}{\partial \Sigma} = 0 = \frac{N\Sigma}{4\pi^2} \left\{ \Sigma^2 \ln \left[1 + \frac{\Lambda^2}{\Sigma^2} \right] - M^2 \ln \left(1 + \frac{\Lambda^2}{M^2} \right) + \Theta(\mu - \Sigma) [2\mu\sqrt{\mu^2 - \Sigma^2} - 2\Sigma^2 \ln((\mu + \sqrt{\mu^2 - \Sigma^2})/\Sigma)] \right\}. \quad (12)$$

Let $\mu < M$ and $M \ll \Lambda$. Then at $\Sigma > \mu$ eq. (12) coincides with stationary equation for the potential $V_0(\Sigma)$ (9). Hence, in this case the single solution of (12) is $\Sigma_1 = M$. At $\Sigma < \mu$ eq. (12) takes the form:

$$\Sigma f_\mu(\Sigma) \equiv \Sigma \left\{ 2\mu\sqrt{\mu^2 - \Sigma^2} - M^2 \ln \left(1 + \frac{\Lambda^2}{M^2} \right) + \Sigma^2 \ln[(\Sigma^2 + \Lambda^2)/(\mu + \sqrt{\mu^2 - \Sigma^2})^2] \right\} = 0. \quad (13)$$

It is easily to show that at $M \ll \Lambda$ the function $f_\mu(\Sigma)$ is monotonically increasing one over variable $\Sigma \in [0, \mu]$. Obviously, that $f_\mu(\mu) < 0$ at $\mu < M$, $f_\mu(\mu) > 0$ at $\mu > M$ and $f_\mu(\mu) = 0$ at $\mu = M$. Consequently, at $\mu < M$ the single solution of eq. (13) is the point $\Sigma_2 = 0$. So, at $\mu < M$ the potential (11) has two stationary points: $\Sigma_1 = M$ and $\Sigma_2 = 0$. At $\Sigma_2 = 0$ there is a local maximum of the potential. But the point $\Sigma_1 = M$ is the point of global minimum for $V_\mu(\Sigma)$. Hence, at $\mu < M$ the chiral symmetry of the NJL model is spontaneously broken and fermions have mass M .

Suppose now that $M < \mu < \mu_{1c}(M)$, where

$$\mu_{1c}(M) = \left[\frac{M^2}{2} \ln \left(1 + \frac{\Lambda^2}{M^2} \right) \right]^{1/2}. \quad (14)$$

It is evident that for such values of the chemical potential $f_\mu(0) < 0$ and $f_\mu(\mu) > 0$. So the monotonically increasing function $f_\mu(\Sigma)$ (remember $M \ll \Lambda$) crosses Σ axis in the single point $\Sigma_3(\mu, M)$, which has properties:

$$\begin{aligned} \Sigma_3(\mu, M) &\rightarrow M & \mu &\rightarrow M \\ \Sigma_3(\mu, M) &\rightarrow 0 & \mu &\rightarrow \mu_{1c}(M). \end{aligned} \quad (15)$$

We have to remark, that in the case under consideration the point $\Sigma_1 = M$ is no more the solution of the eq. (12), but $\Sigma_2 = 0$ is the local maximum of the potential (11) as before. So at $M < \mu < \mu_{1c}(M)$ there is a spontaneous breaking of the symmetry in the NJL model, and fermions acquire mass $\Sigma_3(\mu, M)$.

At $\mu > \mu_{1c}(M)$ eqs. (12), (13) have the solution $\Sigma_2 = 0$ only. So at $\mu > \mu_{1c}(M)$ we have massless phase A of the theory. Here the chiral symmetry is not broken. In the critical point $\mu_{1c}(M)$ there is a phase transition of the second order from the massive phase to the massless one, because the order parameter – fermionic mass – is a continuous function, when $\mu = \mu_{1c}(M)$ (see (15)).

Now we are going to prove that indeed there are two different massive phases of the theory. The first one – the phase B – is situated, when $\mu < M$ and the second one – the phase C – is realized, when $M < \mu < \mu_{1c}(M)$. So we shall try to investigate the thermodynamic potential (TP) $\Omega(\mu)$ of the NJL system at the point $\mu = \mu_{2c}(M) \equiv M$. Let

$$\Omega(\mu) \equiv \begin{cases} \Omega_B(\mu), & \mu < M \\ \Omega_C(\mu), & M < \mu < \mu_{1c}(M). \end{cases} \quad (16)$$

Recall, that TP is the value of the effective potential at the point of global minimum. Hence, $\Omega_B(\mu) = V_\mu(M)$, $\Omega_C(\mu) = V_\mu(\Sigma_3(\mu, M))$. Now let us calculate derivatives of $\Omega_B(\mu)$ and $\Omega_C(\mu)$ at the point $\mu = M$. First of all one should note that from (15) it follows $\Omega_B(M) = \Omega_C(M)$. From (9-11) we have

$$\begin{aligned} \Omega_B(\mu) = V_\mu(M) = V_0(M) = \frac{N}{16\pi^2} \{ M^4 - \\ - M^4 \ln(1 + \Lambda^2/M^2) - \Lambda^4 \ln(1 + M^2/\Lambda^2) \}. \end{aligned} \quad (17)$$

It is clear, that $\Omega_B(\mu)$ is μ independent function, that is why all its derivatives over μ identically equal to zero in the region $\mu < M$, including the case $\mu \rightarrow M_-$. The first derivative of the $\Omega_c(\mu)$ is:

$$\frac{d\Omega_c(\mu)}{d\mu} = \left\{ \frac{\partial V_\mu(\Sigma)}{\partial \mu} + \frac{\partial V_\mu(\Sigma)}{\partial \Sigma} \frac{\partial \Sigma}{\partial \mu} \right\} \Big|_{\Sigma=\Sigma_3(\mu, M)}. \quad (18)$$

Since Σ_3 is the solution of the stationary equation (12), the second item in (18) equals to zero. Taking into account (11), we have:

$$\frac{d\Omega_c(\mu)}{d\mu} = \left. \frac{\partial V_\mu(\Sigma)}{\partial \mu} \right|_{\Sigma=\Sigma_3} = -\frac{N}{3\pi^2}(\mu^2 - \Sigma_3^2)^{3/2}. \quad (19)$$

According to the eq. (15) expression (18) equals to zero, when $\mu \rightarrow M_+$. So the first derivative of the $\Omega(\mu)$ is continuous function at the point $\mu = M$. In the following we need the relations, which are consequences of eq. (13):

$$\begin{aligned} \frac{d\Sigma_3}{d\mu} \equiv \Sigma'_3 &= \left\{ \frac{\partial f_\mu(\Sigma)}{\partial \mu} \left[\frac{\partial f_\mu(\Sigma)}{\partial \Sigma} \right]^{-1} \right\} \bigg|_{\Sigma=\Sigma_3} = \\ &= -\frac{2\sqrt{\mu^2 - \Sigma_3^2}}{\Sigma_3 \{ \ln[(\Sigma_3^2 + \Lambda^2)/(\mu + \sqrt{\mu^2 - \Sigma_3^2})^2] - \Lambda^2/(\Sigma_3^2 + \Lambda^2) \}}, \end{aligned} \quad (20)$$

$$\frac{d^2\Sigma_3}{(d\mu)^2} \equiv \Sigma''_3 = -\frac{2(\mu - \Sigma_3\Sigma'_3)}{\Sigma_4\sqrt{\mu^2 - \Sigma_3^2}\{\dots\}} + O(\mu^2 - \Sigma_3^2). \quad (21)$$

Here in the braces in (21) one should mean the same expression as in the braces in eq. (20). Evidently, that $\Sigma'_3 \rightarrow 0$ at $\mu \rightarrow M$, but the second derivative Σ''_3 at $\mu \rightarrow M$ turns into $(-\infty)$. Let us now find second and third derivatives of the TP $\Omega_c(\mu)$. Taking into account eqs. (20-21) and (19), we obtain

$$\frac{d^2\Omega_c(\mu)}{(d\mu)^2} = -\frac{N}{\pi^2}(\mu^2 - \Sigma_3^2)^{1/2}(\mu - \Sigma_3\Sigma'_3) \quad (22)$$

$$\begin{aligned} \frac{d^3\Omega_c(\mu)}{(d\mu)^3} &= -\frac{N}{\pi^2} \frac{(\mu - \Sigma_3\Sigma'_3)^2}{\sqrt{\mu^2 - \Sigma_3^2}} - \frac{N}{\pi^2} \sqrt{\mu^2 - \Sigma_3^2} (1 - \\ &- (\Sigma'_3)^2 - \Sigma_3\Sigma''_3). \end{aligned} \quad (23)$$

Expression (22) at $\mu \rightarrow M$ turns into zero, but the third derivative of the TP $\Omega_c(\mu)$ at $\mu \rightarrow M$ turns into infinity. Hence, at the point $\mu_{2c} = M$ we have discontinuous third derivative of the thermodynamic potential $\Omega(\mu)$. According to the criterion of the phase transitions [9], this mean that at the point $\mu_{2c}(M)$ we have phase transition of the second order from massive

phase B (at $\mu < M$) to the massive phase (at $M < \mu < \mu_{1c}(M)$). The main physical quantity, which takes different values in those phases, is the density of particles number:

$$n = -\partial\Omega(\mu)/\partial\mu. \quad (24)$$

With the help of eqs. (24) and (17) one can easily show that in the phase the particles density $n_B \equiv 0$. But in the phase (see (19)) we have:

$$n_c = \frac{N}{3\pi^2}(\mu^2 - \Sigma_3^2(\mu, M))^{3/2} \neq 0. \quad (25)$$

Resuming, we should note that in quantum systems with NJL Lagrangians at nonzero chemical potentials there are three different phases. The first one is massless chirally symmetric phase A, the second B and the third C are massive with spontaneous breaking of chiral symmetry phases. The new phase C is not observed in low dimensional four - fermionic models [10, 11].

In the present paper we have restricted ourselves by condition $M \ll \Lambda$. But in the forthcoming article by Vshivtsev, Zhukovsky and Klimenko the whole range of variations for the parameter M is considered. In this case the same phases A,B and C of the NJL model are possible. But the critical behaviour of the model is more various. Indeed, we have observed on the phase diagram μ, M of the model several tricritical points as well, as the phase transitions of second and first orders.

References

- [1] A.S. Vshivtsev, K.G. Klimenko, Pis'ma Zh. Eksp. Teor. Fiz. **64**, 313 (1996).
- [2] Y. Nambu and G. Jona-Lasinio, Phys.Rev. **122**, 345 (1961).
- [3] V.G. Vaks, A.I. Larkin, Zh. Eksp. Teor. Fiz. **40**, 282; 1392 (1961) (in russian); B.A. Arbuzov, A.N. Tavkhelidze, R.N. Faustov, Doklady Akademii Nauk USSR **139**, 345 (1961) (in russian).
- [4] S. Kawati and H. Miyata, Phys.Rev. **D23**, 3010 (1981).

- [5] J. Fuchs, Z.Phys. **C22**, 83 (1984); V. Bernard, U.-G. Meissner and I. Zahed, Phys.Rev. **D36**, 819 (1987); Chr.V. Christov and K. Goeke, Acta Phys.Pol. **B22**, 187 (1991); D. Ebert, Yu.L. Kalinovsky, L. Münchow and M.K. Volkov, Int.J.Mod.Phys. **A8**, 1295 (1993).
- [6] D. Ebert and M.K. Volkov, Phys.Lett. **B272**, 86 (1991); S.P. Klevansky and R.H. Lemmer, Phys.Rev. **D39**, 3478 (1991); M. Faber, A.N. Ivanov, M. Nagy and N.I. Troitskaya, Mod.Phys.Lett. **A8**, 335 (1993).
- [7] T. Inagaki, T. Muta and S.D. Odintsov, Mod.Phys.Lett. **A8**, 2117 (1993); E. Elizalde, S. Leseduarte and S.D. Odintsov, Phys.Rev. **D49**, 5551 (1994); Phys.Lett. **B347**, 33 (1995); D.K. Kim and I.G. Koh, Phys.Rev. **D51**, 4573 (1995).
- [8] L. Dolan and R. Jackiw, Phys.Rev. **D9**, 3320 (1974).
- [9] I.P. Bazarov, E.V. Gevorkyan, P.N. Nikolaev, *Thermodynamics and Statistical Physics*, Moscow State Univ. Press, Moscow (1986).
- [10] V.A. Osipov, V.K. Fedyanin, Teor. Mat. Fiz. **73**, 393 (1987) (in russian); K.G. Klimenko, Teor. Mat. Fiz. **75**, 226 (1988) (in russian).
- [11] K.G. Klimenko, Z.Phys. **C37**, 457 (1988).